

14.4: Directional Derivatives and the Gradient Vector

Calculus III

College of the Atlantic

1. Some warm-ups/review.

(a) Let $f(x, y) = 3x^2y + \sin(xy)$.

i. Calculate the differential of $f(x, y)$.

ii. Evaluate the differential of $f(x, y)$ at $(1, 2)$.

(b) Let $g(x) = x^2$. Use difference quotients with $\Delta x = 0.1$ to estimate $g'(2)$.

2. Let $f(x, y) = 3x^2y$.

(a) Find the gradient of f at $x = 1, y = 2$.

(b) Find the directional derivative $f_{\vec{u}}$ at $(1, 2)$ for the following \vec{u} 's:

i. $\vec{u} = \vec{i}$

ii. $\vec{u} = \vec{j}$

iii. $\vec{u} = \vec{i} + \vec{j}$

iv. $\vec{u} = \vec{i} - \vec{j}$

v. $\vec{u} = -\vec{i}$

vi. $\vec{u} = -\vec{j}$

(c) In what direction \vec{u} is $f_{\vec{u}}$ the largest?

(d) In what direction \vec{u} is $f_{\vec{u}}$ the smallest?

(e) In what direction \vec{u} is $f_{\vec{u}}$ the zero?

3. Consider the function $f(x, y) = x^2 + 4y^2$.

(a) Sketch contour lines for the function in the first quadrant.

(b) Calculate the gradient vector for general x, y .

(c) Determine the value of the gradient vector at the following points:

i. $(1, 1)$

ii. $(1, 2)$

iii. $(2, 1)$

iv. $(2, 2)$

(d) Draw the above gradient vectors on your contour plot sketch. Do the values make sense geometrically?

(e) What is the rate of change of f at $(2, 2)$ in the direction $\vec{u} = -\vec{i} + 2\vec{j}$?

(f) In what direction is the rate of change of f at $(2, 2)$ the largest? I.e., in what direction is the function the steepest uphill?

(g) In what direction is the rate of change of f at $(2, 2)$ the smallest? I.e., in what direction is the function the steepest downhill?

(h) In what direction is the rate of change of f at $(2, 2)$ zero? I.e., in what direction does the function not change?

4. A caterpillar is on a metal surface whose temperature is given by $T(x, y) = 3x^2y - y^3$. The caterpillar does not like heat. It is at the point $(5, 1)$.
- In what direction should it move so that it gets cooler as quickly as possible?
 - If it initially moves at 0.8 cm/s, at what rate does the caterpillar experience a temperature decrease?
5. A bird is flying through a large cloud of pollution whose distribution is given by $\rho(x, y, z) = xz + 3x^2y - y^3$ in units of grams per cubic meter, where x , y , and z are measured in miles. The bird does not like pollution. It is at the point $(1, 2, 1)$.
- In what direction should it move so that it gets to cleaner air as quickly as possible?
 - What are the units of the gradient vector?
 - If it initially flies at 1.2 m/s, at what rate does the bird experience a pollution decrease?
6. Consider the function $f(x, y, z) = e^{-(x^2+y^2+z^2)}$.
- Calculate $\vec{\nabla}f$.
 - Determine the gradient vector at the following points
 - $(0, 0, 0)$
 - $(1, 0, 0)$
 - $(0, 0, 1)$
 - $(1, 1, 1)$
 - What is the gradient vector at the origin? What does your answer mean?
 - What is the directional derivative in the $-\hat{z}$ direction at the point $(1, 0, 0)$.
 - What is the directional derivative in the $-\hat{z}$ direction at the point $(0, 0, 1)$.